

EFFECTS OF VARIETIES ON THE GROWTH AND YIELD OF SOME TOMATOES (*Solanum esculentum* L.) IN MUBI, NORTHERN GUINEA SAVANNAH, NIGERIA

I. D. Bake

Corresponding Author's Email: ibrahimdaudabake@gmail.com

Department of Crop Science
Adamawa State University, P.M.B 25 Mubi, Nigeria

ABSTRACT

The study determined the effect of varieties on the growth and yield of some seeds in Mubi Northern Guinea Savannah Nigeria. Tomatoes which were obtained from Mubi main market. The experimental treatments consist of three different varieties of tomatoes viz; (Seria, UC, and UTC) respectively. The treatments were laid out in a Randomized Complete Block Design (RCBD) with three replications. The plots size was measured 3m x 3m with alley of 1m between replication and 1m between plots. Tomatoes seed were sown directly into the soil by dibbling method Sowing was done by hand using hoe and after germination the seedlings were thinned to one per stand being a set required population for better yield. The results of varieties on plant height showed a significant difference ($P \leq 0.01$) level of significance at 3, 6, and 9 WAS, UTC recorded the highest mean value (23.17) followed by Seria (14.47) with UC recording the least (9.76) at 3 WAS. Similarly, at 6 and 9 WAS UTC recorded the highest mean value of (60.69) and (78.52) followed by Seria at (78.08cm) at 9 WAS and (49.75cm) at 6 WAS where UC recorded the least at (41.02cm) and (61.16cm) at 6 and 9 WAS. The results obtained agree with the findings (Iqtidar et al, 2006 and Keskar et al; 2011). The number of branches per plant at 3, 6 and 9 WAS, the results showed a significant of ($P \geq 0.01$) differences at 3,6 and 9 WAS. UTC recorded the highest mean value (20.26), (14.66) seria at 9 WAS followed by UC (14.80) and (9.20) seria at 6 WAS.

Keywords: Growth, Varieties, Okra, and yield.

INTRODUCTION

Foolad (2018), found that, tomato is the sixth most valuable crop in the world worth US\$ 87.9 billion in 2021 and is grown in all soil types on a small scale for family use and commercial purposes. Tomato belongs to the solanaceae family and the genus *Solanum* which is a small genus within the large and diverse genera that consists approximately 90 genera (Singh, 2019) and is one of the most consumed vegetables in the world. Tomatoes can be eaten fresh or in processed forms such as sau U, ketchup, puree, paste, powder and soup (Battistuzzi, 2012). Tomato is also a crop which is globally grown, some in the open field others in the green house. Indeterminate varieties are usually grown in greenhouses (Stoleru, 2020). Tomatoes fruits exist in many shapes appearing large and round, oval or elongated, depending on the variety. The fruits may be orange, yellow or red when ripe, usually with numerous kidneys or peer shaped, hairy, light brown seeds (Tindall, 2018). Tomato is a warm season plant that requires temperature of about 20-27°C for most cultivars. Excessive rainfall and high relative humidity can be harmful to it. A deep loamy soil well drained rich in organic material with a pH of 6.2 – 6.8 for optimum growth and development. It is a nutritious vegetable with increasing demand in Nigeria. As documented by United States Dietary Allowance (2008), tomato fruit provide 95% water of its edible portion and the remaining 5% compose of miscellaneous compound among which include carotenoids, ascorbic acids, alcohol, insoluble solids (proteins) cellulose, pectins, polysaccharides and inorganic compound which give the fruit its characteristic flavor and aroma. In addition, (Stoleru, 2020) revealed that tomatoes are among the food plants to which moderate level of anti-cancer activities is associated. It also contains Vitamin A which is a remedy for night blindness. The world's cultivation of tomato is put

to about 4 million hectares of which 63,482 hectares are estimated to be cultivated. In Nigeria. FAO (2010) reported an estimated annual production of 1.7 million tones. But however, this was still far below demand as there was an increasing demand from human consumption. However, most of the deficits were offset by importation. Although, despite a drive towards increased tomatoes production in Nigeria, an optimum production has not been realized by the framers to meet its demand among consumers. Generally, tomato production in Nigeria has been low because of low yields obtained by small scale farmers resulting from the use of unimproved local varieties with low yielding capacity which often grown in mixtures and this may prevent exploitation of crop productive capacity, environmental hazards such as drought and incidence of pest and diseases, plant population including soil fertility decline and a host of other factors have also kept tomato yield output relatively low (Adekiya and Ojeniyi, 2012). Therefore, effect of varietal influence on the growth and yield of three tomatoe varieties need to be evaluated to replace the low yielding local varieties currently in use by farmers. Tomato productivity in Nigeria is rated very low at 1.5 to 14 tons per hectare compared to world average of 56 tons per hectare (FAO, 2014). The most common limiting factors to tomato productivity are pests and diseases, drought, soil erosion, and declining soil fertility across Nigeria (Birungi, 2020). Cultural practices associated with tomato growth and fruit developed have been investigated, but the findings are not conclusive. The situation is worst in developing countries such as Nigeria, where there is insufficient information on the cultural practice of some varieties of tomatoes. "Tomato (*Solanum lycopersicum*) is a widely cultivated crop, and its yield and quality are significantly influenced by varietal differences. Despite the importance tomato production, there is limited research on the comparative growth and yield performance tomato varieties under the same environmental conditions. This knowledge gap hinders the ability and gardeners to make informed decisions about which varieties to cultivate, leading to reduced yields and decreased productivity. For this reason, it is difficult to define the optimal development condition to maximize the production of tomatoes. To have more information, it would be of interest to give an insight on how agronomic and environmental factors affect some varieties. The research therefore be conducted to investigate the influence of varietal effect on the growth and yield of three tomatoe varieties.

Justification of the Study

The production of tomato could not meet up with the demand for food, industrial uses and export due to lack of proper cultural practices. There is a need to avail to the famers the best variety of tomatoes that will performance well. Tomato varieties exhibit distinct characteristics, and understanding their growth and yield of different tomato varieties under the same environmental conditions. So, by investigating the effect of varietal influence on the growth and yield of three tomato varieties, this will study will aim to address a significant knowledge gap, providing practical and scientific benefits that can impact tomato production and food security.

Objective of the study

- i. evaluate the effect of varieties on the growth and yield of tomatoes (*Solanum esculentum* L.)
- ii. determine the best varieties of tomatoes (*Solanum esculentum* L.) that can give maximum yield in study area.

MATERIALS AND METHODS

Field experiment was conducted at the Department of crop science, teaching and research farm, food and agricultural organization/tree crop programme (FAO/TCP) farm Faculty of Agriculture, Adamawa State University Mubi, Nigeria. Mubi located in the northern guinea savanna of Nigeria is situated

between latitude $10^{\circ}11'$ and $10^{\circ}30'$ north of the equator and between longitude $13^{\circ}10'$ and $13^{\circ}30'$ east of the Greenwich meridian and at altitude of 969m above mean sea level (MSL). The mean rainfall is 900mm and a minimum temperature of 18°C harmattan period and 40°C as maximum in April (Adebayo, 2020).

Treatment and Experimental Design

The experiment design was laid out in a Randomized Completed Block Design (RCBD). The study was carried out in Mubi North Local Government Area. The treatments consists of three (3) tomato varieties viz: T1 (seria), T2 (UC) and T3 (UTC). The treatments were laid out in a Complete Block Design (RCBD) in three (3) replicates

Sources of Tomato Seed

Seria, UC and UTC seeds was bought from Mubi local market.

**Table 1: Physio-Chemical Properties of the Experimental site before Cropping
 Soil Physical Properties**

	Sand	Silt	Clay	Textural classes	B.D g/cm ³	P.D g/cm ³	Porosity (%)
	%						
A	68.8	14.4	16.8	Sandy Loam	1.49	2.384	37.5
B	70.8	11.4	17.8	Sandy Loam	1.46	2.43	39.9177

Soil Chemical Properties

	pH(1:2)	EC(ds/m)	Org.C(%)	Org.M(%)	TN(%)	Av-P(mg/kg)	Ca	Mg	Na	K	TEB	H	Al	TEA	ECEC	PBS(%)	ESP(%)
A	6.7	0.070	1.057	1.823	0.099	9.260	1.849	2.249	0.0883	0.555	4.736	0.576	1.508	1.508	6.244	85.672	1.322
B	6.80	0.09	0.54	0.93	0.05	8.74	8.80	1.20	0.78	0.23	11.01	0.74	1.13	1.87	12.88	88.72	6.09

Nursery Operation of Tomatoes

Two nursery beds were raised 45cm × 45cm each, separated 1cm apart. 1kg of poultry dropping was spread on each beds and allowed to decompose for 14days. The seeds of the three varieties were sown by broadcasting method on each of the raised seed bed. All nursery management activities were strictly adhered to and the nursery duration took four weeks, before transplanting the tomatoe seedlings on the research field.

Transplanting of Tomatoes Seedlings

The seedlings of the three varieties tomatoes were transplanted four weeks after broadcasting on already ploughed and harrowed. Transplanting was done early in the morning after a light shower of rainfall using the spacing of 60cm × 60cm.

Cultural Practices

Pest and Disease Control of Tomatoes

In control possible incidence of pest on disease associated with tomato, insecticide such as cypermethrine was applied at growth stages on weekly bases

Weed control

Hand weeding using hoe was done at three and six weeks after transplanting to control weeds competition for space, nutrient and sunlight required for photosynthesis.

Data Collection

Data were collected on the following parameters;

Plant height (cm): Five plants were tagged. Height of the tagged plants were measured each from the base to tip of the stem using a meter scale at harvest. Number of branches: total number of primary branches of selected and tagged plants (five plant) in the treatment were counted at the time of last harvest or picking of fruits and the mean was recorded. Stem girth (mm): total number of primary stem of selected tagged plants under a treatment was counted at the time of last harvest or picking of fruits and the mean value was evaluated by selecting some plants randomly from treatment and at the time of last harvest. This was done by measuring the thickness of the stem at harvest.

Days to 1st flowering: The number of days taken from date of sowing to date at which 1st of the plant population flowered was noted and then the average was determined.

Days to 50% flowering: this is the period whereby 50% of the plant produce flowers

Fresh Fruits weight (FFW): The edible fruits weight in kg at tagged plants. Each treatment was recorded with the help of weighing balance. The total weight of fruits was divided from the tagged plants the number of fruits to obtain average fresh fruits weight by measuring the fruits weight of each plant in kg.

Statistical Analysis

Data obtained was subjected to the analysis of variance (ANOVA) and treatment mean was compared using the Ducan Mutiple Range Test (DMRT) at 5%.

Results and Discussion

Table .1 Effect of Varieties on Plants Height(cm), and Number of Branches at 3, 6 and 9 WAS in 2024 Cropping Season

Treatments	PH3	PH6	PH9	NB3	NB6	NB9
Seria	14.47 ^b	49.75 ^b	76.08 ^{ab}	4.93 ^b	9.20 ^b	14.66 ^b
UC	9.76 ^c	41.02 ^{bc}	61.16 ^b	3.46 ^c	14.80 ^a	11.66 ^c
UTC	23.17 ^a	60.69 ^a	78.57 ^a	7.20 ^a	6.33 ^c	20.26 ^a
P of F	0.011 ^{**}	0.008 ^{**}	0.003 ^{**}	0.041 [*]	0.002 ^{**}	0.008 ^{**}
SE	2.98	4.03	3.17	1.11	1.33	1.177

Keys: PH3: plant height at week 3, PH6: plant height at week 6, PH9: plant height at week 9, NB3: Number of branches at 3 weeks, NB6: Number of branches at 6 weeks, NB9: Number of branches at 9 weeks

The results of varieties on plant height at 3, 6 and 9 weeks are presented on Table .1 above showed a significant difference ($P \leq 0.01$) level of significance at 3, 6, and 9 WAS, UTC recorded the highest mean value (23.17) followed by seria (14.47) UC recorded the least at (9.76) at 3 WAS. Similarly, at 6 and 9 WAS UTC recorded the highest mean value of (60.69) and (78.52) followed by Seria at (78.08cm) at 9 WAS and (49.75cm) at 6 WAS UC was recorded the least at (41.02cm) and (61.16cm) at 6 and 9 WAS. The plants number of branches at 3,6 and 9 WAS (Table .1). showed a significant ($P \geq 0.01$) differences at 3,4 and 9 WAS. UTC recorded a high mean value (20.26), (14.66) seria at 9 WAS followed by UC (14.80) and (9.20) seria at 6 WAS UC recorded the least with (3.46), respectively.

The results of the effect of varieties on the plant stem girth at 3,6, and 9 WAS showed that at 9 WAS Seria recorded the high mean value of (354.42 mm/plant) followed by UTC at (11.97 mm/plant) at 3 WAS, and (9.93 mm/plant) at 6 WAS while at 3,6 and 9 WAS UC recorded the least mean value at 3 WAS at (4.00 mm/plant) at ($p \geq 0.01$) level of significance (Table at .2). The effect of varieties on days to first and 50% flowering shauce recorded that UC has the high mean value at DFF with (34.66) and (42.00) at DFPP and UTC recorded the least with (24.00) DFF and (28.33) DFPP with ($P \geq 0.01$) level of significance.

Table 4.2 Effect of Varieties on Stem Girth (mm), Days to First and Fifty (50) % Flowering in 2024 Cropping Season

Treatments	SG3	SG6	SG9	DFF	DFFP
Seria	4.86 ^b	8.06 ^b	12.42 ^a	24.66 ^b	28.66 ^b
UC	4.00 ^{bc}	6.39 ^c	7.71 ^c	34.66 ^a	42.00 ^a
UTC	12.34 ^a	9.93 ^a	11.97 ^b	24.00 ^{bc}	28.33 ^{bc}
P of F	0.348 ^{NS}	0.009 ^{**}	0.422 ^{NS}	0.001 ^{**}	0.012 ^{**}
SE	5.76	0.73	281.52	1.61	3.45

Keys: SG3: Stem Girth at 3 weeks, SG6: Stem Girth at 6 weeks, SG9: Stem Girth at 9 weeks, DFF: Days to first flowering, DFFP: Days to fifty percent flowering

Table 4.3 Effect of Varieties on Fruit Yield (Kg/ha⁻¹) in 2024 Cropping Season

Treatment	FY1	FY2	FY3	FY4	FY5	FY6	FY7
Seria	282.20 ^a	638.53 ^a	164.53 ^a	233.60 ^a	203.60 ^a	84.50 ^c	96.93 ^c
UC	120.00 ^b	148.66 ^{bc}	136.86 ^b	131.11 ^c	162.46 ^c	134.33 ^b	160.73 ^b
UTC	242.53 ^{ab}	162.46 ^b	135.66 ^b	188.53 ^b	196.13 ^b	147.13 ^a	208.26 ^a
P of F	0.145 ^{NS}	0.362 ^{NS}	0.587	0.001	0.004	0.371	0.008
SE	72.71	358.88	30.27	13.60	7.63	43.18	23.05

FY: Fruit Yield(Kg/ha⁻¹)

The results of the effect of varieties on the fruits yield at FY2 and FY1, seria recorded the highest mean value of (638.53) and (282.20) respectively followed by UTC at (242.53) FY1 UC recorded the lowest at FY1 with 120.00 fruits yield.

Effect of Varieties on Plants Height, and Number of Branches at 3, 6 and 9 WAS

The result showed a significant difference ($P \leq 0.01$) level of significance at 3, 6, and 9 WAS, UTC recorded the highest mean value (23.17) followed by Seria (14.47) where UC recorded the least (9.76) at 3 WAS. Similarly, at 6 and 9 WAS UTC recorded the highest mean value of (60.69) and (78.52) followed by Seria at (78.08cm) at 9 WAS and (49.75cm) at 6 WAS with UC recording the least at (41.02cm) and (61.16cm) at 6 and 9 WAS. The results obtained agrees with the findings at (Iqtidar *et al*, 2006 and Keskar *et al*; 2011). The plants number of branches showed a significant ($P \geq 0.01$) differences at 3,4 and 9 WAS. UTC recorded with the highest mean value of (20.26), (14.66) seria at 9 WAS followed by UC (14.80) and (9.20) seria at 6 WAS UC recorded the least with (3.46), respectively. The results obtained are in tandem with those of Singh (2020), Adekiya (2017) and Agbede (2008) who reported that application of poultry manure and the use of UTC tomato varieties gave the highest number of flowers per plant.

Effect of Varieties on Stem Girth, Days to First and Fifty (50) % Flowering

The results of the effect of varieties on the plant stem girth at 3,6, and 9 WAS showed that at 9 WAS Seria recorded the highest mean value of (354.42 mm/plant) followed by UTC at (11.97 mm/plant) at 3 WAS, and (9.93 mm/plant) at 6 WAS while at 3,6 and 9 WAS UC recorded the least mean value at 3 WAS with (4.00 mm/plant) at ($p \geq 0.01$) level of significance (Table at 4.2) This observation was corroborated with the findings of (Bache and (Helathcote, 1996, Adediran, 2003), that both UC, UTC and Seria increased. The effect of varieties on days to first and 50% flowering recorded that UC has highest mean value at DFF with (34.66) and (42.00) at DFPF and UTC recorded the least mean of (24.00) DFF and (28.33) DFPF with ($P \geq 0.01$) level of significance.

Effect of Varieties on Fruit Yield

The results of the effect of varieties on the fruits yield at FY2 and FY1, seria recorded the highest mean value of (638.53) and (282.20) respectively followed by UTC at (242.53) FY1 where UC recorded the lowest at FY1 with 120.00 fruits yield. Highly significant difference was recorded on leaf area on FY2. This results agrees with the findings of the following researchers Ramachandra and Thimmaraju 1983; Ehigiator, 1990; Singh and Whitehead, 1996; Egharevba and Ogbe, 2002 Olaniyi, 2007; Jombo *et al*; 2012). They also reported on the response of *Amaranthus* to different varieties and rates of fertilizer on fresh and dry weight, leaf area index, absolute crop growth rate and fresh yield of *Amarathus*.

SUMMARY

The study determines the effect of varieties on the growth and yield of some tomatoes in Mubi Northern Guinea Savannah Nigeria. Seeds of tomatoes were obtained from Mubi main market. The experimental treatments consisted of three different varieties of tomatoes viz; (Seria, UC, and UTC) respectively. The treatments were laid out in a Randomized Complete Block Design (RCBD) with three replications. The plots size was measured 3m x 3m with alley of 1m between replication and 1m between plots. Tomatoe seeds were sown directly to the soil by dibbling method Sowing was done by hand using hoe and after germination the seedlings were thinned to one per stand to set required population for better yield. The results of varieties on plant height showed a significant difference ($P \leq 0.01$) level of significance at 3, 6, and 9 WAS, UTC recorded the highest mean value (23.17) followed by Seria (14.47) where UC recorded the least at (9.76) at 3 WAS. Similarly, at 6 and 9 WAS UTC recorded the highest mean value of (60.69) and (78.52) followed by Seria at (78.08cm) at 9 WAS and (49.75cm) at 6 WAS with UC recording the least at (41.02cm) and (61.16cm) at 6 and 9 WAS. The results obtained agree with the findings of (Iqtidar *et al*, 2006 and Keskar *et al*; 2011). Results of plants number of branches at 3,6 and 9 WAS showed at the results showed a significant of ($P \geq 0.01$) differences at 3,4 and 9 WAS. UTC recorded the highest mean value (20.26), (14.66) seria at 9 WAS followed by UC (14.80) and (9.20) seria at 6 WAS

CONCLUSION

Based on the above findings of experiment, it can be concluded that the use of Seria, UC and UTC that recorded significant differences between the different varieties on the yield of tomatoes and even their combination can be beneficial to farmers. It is therefore recommended that farmers may opt for either of the three varieties of tomatoes for a better productivity.

RECOMMENDATIONS

Based on the findings of this study, it may be recommended that it would be advisable to farmers to use either of the three varieties of tomatoes so that better yields may be achieved with moderate application of fertilizer. This should be encouraged to forestall vegetative growth at the expense of seed formation and for maximum growth and yield of tomato in the study area.

REFERENCES

- Acquaah, E. K. (2002). Seeing below the surface: making soil processes visible to Ugandan smallholder farmers through a constructive and experiential extension approach. *Agriculture and Human Values*, 35(2), 425-440.
- Adebayo, A.A. Adebayo, A.A and Tukur A.L. (2020). Climate ii. Rainfall in *Adamawa State in Maps*. (eds) paraclete publishers Yola. Pp. 23-26
- Adeniyani, O. N. and S. O. Ojeniyi. (2005). Effect of poultry manure, NPK 15-15- 15 and combination of their reduced levels on maize growth and soil chemical properties. *Nigerian Journal of Soil Science* 15:34-41.
- Agbede, T.M., S.O. Ojeniyi and A.J. Adeyemo, (2012). Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in southwest, *Nigeria. Am.-Eurasian J. Sustain. Agric.*, 2: 72-77.
- Angole, S. (2010), "Tobacco to tomatoes: A phylogenetic perspective on fruit diversity in the Solanaceae", *Journal of Experimental Botany*, Vol. 53/377, pp. 2001-2022.
- Asadu L.A and B.O. Unagwu (2012). Effect of Combined Poultry manure, and Inorganic fertilizer on maize performance in an Ultisol of Southeastern Nigeria. In: *Nigerian Journal of Soil sciences*. Vol 22(2) Pp 79-87.
- Battistuzzi, M. Á. G. and Valenciano, J. D. P., (2012). Analytical model for the global consumption of tomatoes-The Spanish case. *African Journal of Agricultural Research*, 7(15), 2328- 2335.
- Birungi, G. & Ngabirano, H. (2020). Pesticide Use in Vegetable Production In Rural Uganda-A Case Study of Kabale District, South Western Uganda.
- Childres, R.T. (2005), "Cytogenetics and evolution", in: Razdan, M. and A.K. Matoo (eds.), *Genetics Improvement of Solanaceous Crops. Vol 2: Tomato*, Science Publishers, New Delhi, India.
- Clinton, M. Q. (1998), "Immunogenicity of papaya mosaic virus-like particles fused to a hepatitis C virus epitope: Evidence for the critical function of multimerization", *Virology*, Vol. 363/1, pp. 59-68.
- FAO (2014) Production-Crops, 2012 data. Food and Agriculture Organization of the United Nations.
- FAO (2010) (Food and Agriculture Organization of the United Nations). FAOSTAT Database. <http://faostat3.fao.org/>
- Foolad, M. R., Ohlson, E. W., & Ashrafi, H., (2018). Identification and mapping of late blight resistance quantitative trait loci in tomato accession PI 163245. *The Plant Genome*, 11(3), 1-16.

- Giovanucci, LA, (1995). Bcarotene and ascorbic acid retention in fresh and processed vegetables. *J. Food Sci.* 64(5):929-936.
- Hartmann, H.T, and D. E KESTER. 2002 Plant propagation principles and practices. 4th edition Englewood cliffs, N.J; Prentice – Hall.
- Mwebaze, F. (2004). Policies to Promote Sustainable Agriculture in Uganda-SWICTH Africa Green.
- Olatunji, S.O., S.A. Ayuba, B.C. Anjembe and S .0 Ojeniyi (2012)- Effect of NPK and Poultry manure on Cowpea and Soil nutrient composition. *In Nigerian Journal of Science* Vol. 108- 113
- Olatunji O and V. U. Oboh (2012) Growth and yield of Okra and Tomato as affected by Pig dung and other manure issue for Economic Consideration in Benue state. In: *Nigerian Journal of Soil Science* Vol. 22 PP 103-107
- Sajjan A. Q., Zhang, L., Zou, D., Sun, Y., Gao, M., and Wang, X. (2002). A comparison of variety, antibiotic resistance genes, and bacterial community in broiler and layer manure following composting. *Environmental Science and Pollution Research*, 1-13.
- Shi, N. and Le Mguer O. V. (2000). Role of plant nutrients in plant growth and physiology. In *Plant Nutrients and Abiotic Stress Tolerance* (pp. 51-93). Springer, Singapore.
- Singh, M., Rajawat, K. S., Ameta, K. D., Kaushik, R. A., Lakhawat, S. S., Shaktawat, and S. K., (2019). Effect of organic manures and chemical fertilizers on fruit characteristics, quality and yield of tomato under naturally ventilated polyhouse. *IJCS*, 7(5), 1881-1884.
- Stoleru, V., Inculet, S. C., Mihalache, G., Cojocar, A., Teliban, G. C., and Caruso, G. (2020). Yield and Nutritional Response of Greenhouse Grown Tomato Cultivars to Sustainable Fertilization and Irrigation Management. *Plants*, 9(8), 1053.
- Tindall HD, (2018). Tomatoes RP. Fruit and vegetable production in warm climates. The Macmillan press Ltd. 13;85.
- Tonny, W.B. (2004). Growth. nutrient uptake and yield Of maize and Okra as influenced by cow dung and plant material and nitrogen fertilizer under different cropping systems. Ph.D. Thesis, University of Ibadan, Nigeria. 22Spp
- Ton, W.D. Sahul M. F. and Togun, A.O. (2004). The influence of maize-stover cow dung and plant material and nitrogen fertilizer on growth. yield and nutrient uptake of amaranth. *Scientia Horticulture* 93: I -8.
- United State Dietary Allowance (USDA). (2008). Food Safety Regulatory Essentials (FSRE) Shelf-Stable, Principles of preservation of shelf-stable dried meat products. United State Department of Agriculture. Food Safety and Inspection Service.
- Wilcox, N. T. Ofor, C. O., and Alozie, L. N. (2003). The Composition of Layer and Broiler Manures, and Implications for Maggot Production and Pond Fertilization. *Waste and Biomass Valorization*, 3(1), 43-48.

Williams, V.M. (2004), "Root-nematode resistance genes in tomato and their potential for future use", *Annual Review of Phytopathology*, Vol. 36, pp. 277-293.

Yahia, F. S. and Brecht, O. N (2012), *Catálogo de Semillas de Tomate*, Banco de Germoplasma de la Universidad Politécnica de Valencia, INIA, Madrid, Spain.